

Wild ginger

Hedychium gardnerianum

What does it look like?

Both species of wild ginger are non-woody perennials, growing up to 2m tall from thick-branching rhizomes (swollen underground stems). Their dense underground root system spread up to one metre deep excluding all other species. Both species produce shiny leaves 20 - 45cm long.

Kahili ginger flowers January to March and produces scented, lemon and yellow flowers with conspicuous red stamens, followed by fruiting spikes with fleshy orange fruits. Yellow ginger produces scented, cream coloured flowers in clusters, from May to June and does not produce fruit.

Both species prefer forest environments, where soils are fertile and damp, thus wild ginger has been termed a 'forest invader'. Areas where it can commonly be seen growing include forests, streamsides, alluvial forest, light gaps and gullies. It cannot tolerate very dry or rocky areas, due to the generally low fertility of the soil in such environments.

Why is it a problem?

Wild ginger can still successfully multiply when rhizomes are damaged. Young plants are very palatable to livestock, and both species tend to invade all areas where stock are excluded, including within native bush and on road sides and riverbanks. Due to its smothering nature, it will eventually replace all other species, being extremely shade tolerant, somewhat tolerant of frost and drought and can withstand immersion in sea water.



Wild ginger

Hedychium gardnerianum

Control Methods

Physical control

Hand pull or dig out small plants and dispose of rhizomes at landfill or burn.

Remove flower heads as a stopgap before spraying or removing the whole plant. This will stop it from seeding that year. Burn the seed heads or dispose of at landfill.

Herbicide control

Cut stems, year round, above pink 'collar' at base and cover cut surface with metsulfuron-methyl (600g/kg active ingredient) at 1g /L of water, or, with metsulfuron gel. Leave stems and leaves on site to rot down. Or,

Foliar spray, year round, with metsulfuron-methyl (600g/kg active ingredient) at 0.5g/L of water plus penetrant at the label rate.

Note: Overspray will kill other (desirable) broadleaf plants and has residual activity in the soil which aids in killing below ground parts. For dense patches keep spray away from roots of vulnerable plants. Don't replant sprayed sites for 6 months. Do not use over or near water bodies or wetlands.

CAUTION: When using any herbicide or pesticide, PLEASE READ THE LABEL THOROUGHLY to ensure that all instructions and safety requirements are followed.

Related Links

Ministry for Primary Industries <http://apps.mpi.govt.nz/applications/nzpests-view/Article/997/Kahili-ginger>

New Zealand Plant Conservation Network http://www.nzpcn.org.nz/flora_search.aspx?scfSubmit=1&scfStart_Res ults=&scfFlora_Category=&scfLatin_Name=Hedychium&scfCommon_Name=&scfFamily_Name=&scfDistrict=&scfAnc istry=&scfFlora_Structural_Class=&scfFlora_Status=

NZ Flora <http://www.nzflora.info/search.html?q=Hedychium>

Unwanted Organisms Register https://www1.maf.govt.nz/uor-cgi/uor.pl/show?species_id=36650

Weedbusters <https://www.weedbusters.org.nz/what-are-weeds/weed-list/wild-ginger/>

Rules

Under Section 52 and 53 of the Biosecurity Act 1993 no person can sell, propagate, breed, distribute or otherwise spread any pest in a Pest Management Plan, or any unwanted organism. Not complying with Section 52 or 53 is an offence under the Act, and may result in penalties noted in Section 157(1).

The National Pest Plant Accord is designed to prevent the sale, distribution and propagation of a set list of pest plants (the Accord list) within New Zealand. If allowed to spread further, these pest plants could seriously damage the New Zealand economy and environment.